

A first-principles derivation of York scaling and the Lichnerowicz–York equation

E. Anderson,^{1,*} J. Barbour,^{2,†} B.Z. Foster,^{3,4,‡} B. Kelleher,^{5,§} and N. Ó Murchadha^{5,¶}

¹*Astronomy Unit, School of Mathematical Sciences,
Queen Mary University of London, London, E1 4NS, UK*

²*The Leibniz Institute at College Farm, South Newington, Banbury, Oxon, OX15 4JG, UK*

³*Institut d’Astrophysique de Paris, 98 bis Bvd. Arago 75014 Paris, France*

⁴*Physics Department, Univ. of Maryland, College Park, Maryland, USA*

⁵*Physics Department, University College Cork, Ireland*

(Dated: February 6, 2008)

The only efficient and robust method of generating consistent initial data in general relativity is the conformal technique initiated by Lichnerowicz and perfected by York. In the spatially compact case, the complete scheme consists of the Arnowitt–Deser–Misner (ADM) Hamiltonian and momentum constraints, the ADM Euler–Lagrange equations, York’s constant-mean-curvature (CMC) condition, and a lapse-fixing equation (LFE) that ensures propagation of the CMC condition by the Euler–Lagrange equations. The Hamiltonian constraint is rewritten as the Lichnerowicz–York equation for the conformal factor ψ of the physical metric $\psi^4 g_{ij}$ given an initial unphysical 3-metric g_{ij} . The CMC condition and LFE introduce a distinguished foliation (definition of simultaneity) on spacetime, and separate scaling laws for the canonical momenta and their trace are used. In this article, we derive all these features in a single package by seeking a gauge theory of geometrodynamics (evolving 3-geometries) invariant under both three-dimensional diffeomorphisms and volume-preserving conformal transformations.

PACS numbers: 04.20.Cv, 04.20.Fy

One traditionally views general relativity (GR) as giving 4-d space-times satisfying the Einstein equations. However, to make easier contact with the rest of physics, one would like to study GR, and gravitation more generally, as a dynamical theory. In the standard way of doing this, due to Arnowitt–Deser–Misner (ADM) [1] and Dirac [2], one introduces some time-function into the space-time. Each time-slice is a three-dimensional Riemannian manifold with a 3-metric g_{ij} . One can view the dynamics of GR as the evolution of these 3-metrics. In this letter, we will restrict our attention to smooth metrics on a fixed manifold that is compact without boundary (CWB).

The 4-d general covariance of the original theory allows free 3-d coordinate transformations on each slice. This means that the configuration space is *superspace*, the space of 3-geometries, identified as the quotient with respect to 3-diffeomorphisms of Riem, the space of all 3-metrics defined on a given CWB manifold. GR in this approach is a theory with constraints. One of the constraints, the *momentum constraint* is a 3-vector condition that just generates the 3-diffeomorphisms. The other constraint, the *Hamiltonian constraint*, is a scalar that represents the freedom one has in choosing the time-function. A 3-metric has six free components; factoring out by the 3-diffeomorphisms and the choice of time-function leaves two degrees-of-freedom per space point, exactly as we expect for the gravitational field. Finding an explicit representation of these two degrees-of-freedom is of major practical and theoretical interest.

Conformal 3-geometries have exactly two degrees-of-freedom per space point. They are obtained by identi-

fying 3-geometries equivalent up to 3-d conformal transformations, $g_{ij} \rightarrow \omega^4 g_{ij}$, where ω is an arbitrary suitably continuous function. One obtains *conformal superspace* (CS) [3], the space of conformal 3-geometries, by quotienting superspace by the conformal transformations.

In the York method [4, 5], one constructs initial data satisfying the constraints in the form that they take on a constant mean curvature (CMC) slice. One picks a 3-metric, a transverse traceless (TT) symmetric two-index tensor on the 3-manifold, and a single number, the value of the CMC. There is a distinguished rescaling of York’s TT tensors under conformal transformations. Its use makes it possible to solve the momentum constraint algebraically, and the Hamiltonian constraint reduces to a well-behaved elliptic scalar equation for the conformal factor ψ , the *Lichnerowicz–York equation* [4, 6]. Because this is to be solved for the conformal factor, the original 3-metric g_{ij} turns out to be unphysical, but the physical 3-metric, $\psi^4 g_{ij}$, is deduced by solving for the conformal factor. One can adjoin to the above constraint system the *lapse-fixing equation* (LFE), another well-behaved elliptic equation. This allows one to construct a CMC foliation of the spacetime and thus generate a 4-manifold satisfying the Einstein equations.

Here, we derive the salient features of the York method from first principles that have not hitherto been used in gravitational theory. We will first explain our principles, define our action, discuss our variational techniques, and derive our results. We conclude by comparing our approach with York’s.

We wish to construct a dynamical theory of 3-geometry

by extremalizing an action defined on a suitable configuration space. Two general principles underlie our construction: 1) Time is derived from change; 2) Motion and size are relative. Principle 1 leads us to construct on our configuration space a theory of ‘timeless geodesics’, parametrized by a freely chosen label λ . Local proper time is emergent in this approach. Principle 2 leads us to build into the theory invariance under λ -dependent 3-diffeomorphisms and conformal transformations. Our principles will allow us to derive York’s approach to GR (in the CWB case) without any prior assumptions of 4-d space-time structure.

Consider the Baierlein–Sharp–Wheeler (BSW) [7] form of the GR action:

$$I_{\text{BSW}} = \int d\lambda \mathcal{L}_{\text{BSW}} = \int d\lambda d^3x \sqrt{g} \sqrt{R} \sqrt{T}, \quad (1)$$

where $g = \det(g_{ij})$, R is the 3-d scalar curvature, and

$$T = G^{ijkl} \frac{dg_{ij}}{d\lambda} \frac{dg_{kl}}{d\lambda} \equiv G^{ijkl} (\dot{g}_{ij} - \mathcal{L}_\xi g_{ij}) (\dot{g}_{kl} - \mathcal{L}_\xi g_{kl}). \quad (2)$$

Here $G^{ijkl} \equiv (g^{ik}g^{jl} - g^{ij}g^{kl})$ is the inverse DeWitt supermetric, and $\mathcal{L}_\xi g_{ij} = \nabla_i \xi_j + \nabla_j \xi_i$ is the Lie derivative of the metric with respect to ξ^i . This action is invariant under local reparametrization of λ , satisfying Principle 1. The emergent proper time t is related to the label λ by $\delta t = N \delta \lambda$, for $N \equiv \frac{1}{2} \sqrt{T/R}$. The covector $\xi_i = g_{ij} \xi^j$ is effectively a gauge auxiliary that renders the action invariant under 3-diffeomorphisms, satisfying the ‘motion is relative’ part of Principle 2.

The BSW action is defined on curves in superspace. We considered generalizations of this action in [8]. As this led us almost uniquely back to the BSW action of GR, it constitutes a new derivation of GR based on our principles. In [9], we extended the BSW action so as to be defined on CS, thus satisfying also the ‘size is relative’ part of Principle 2. One could hope that this move would lead to GR in the York picture—in fact, this is almost correct. While this leads to a self-consistent theory of gravity, it is not GR. Here, we shall instead extend the BSW action to include invariance under conformal transformations *that preserve the global volume* of the 3-metric. This volume-preserving condition allows us to recover standard GR in the York picture.

We will denote a volume-preserving conformal transformation (VPCT) with a ‘hat’:

$$g_{ij}(x) \rightarrow \hat{\omega}(x)^4 g_{ij}(x). \quad (3)$$

One can construct a VPCT $\hat{\omega}$ from any unrestricted conformal transformation ω :

$$\hat{\omega} = \frac{\omega}{\langle \omega^6 \rangle^{1/6}}, \quad (4)$$

where

$$\langle F \rangle = \frac{\int d^3x \sqrt{g} F}{\int d^3x \sqrt{g}}, \quad (5)$$

denotes the global average of some function F . One can express any VPCT $\hat{\omega}$ in this manner.

We implement the conformal symmetry in the BSW action (1) by introducing an auxiliary scalar field ϕ , constructing from it in the manner above the field $\hat{\phi}$. Under a VPCT (4), we declare that $\hat{\phi}$ transforms as

$$\hat{\phi} \rightarrow \frac{\hat{\phi}}{\hat{\omega}}. \quad (6)$$

Consequently, the ‘corrected coordinates’ $\bar{g}_{ij} \equiv \hat{\phi}^4 g_{ij}$ are invariant under a VPCT. We then re-express the BSW action in terms of the variables $\bar{g}_{ij}$, obtaining an action functional of the variables g_{ij} and ϕ . We emphasize that we vary ϕ freely, not $\hat{\phi}$.

On the introduction of the auxiliary variable ϕ , the constituent parts of the BSW action become

$$\begin{aligned} R &\rightarrow \hat{\phi}^{-4} \left(R - 8 \frac{\nabla^2 \hat{\phi}}{\hat{\phi}} \right), \\ T &\rightarrow \hat{T} = \hat{\phi}^{-8} G^{ijkl} \frac{d\hat{\phi}^4 g_{ij}}{d\lambda} \frac{d\hat{\phi}^4 g_{kl}}{d\lambda}, \end{aligned} \quad (7)$$

with

$$\frac{d\hat{\phi}^4 g_{ij}}{d\lambda} = \hat{\phi}^4 \left[\dot{g}_{ij} - \mathcal{L}_\xi g_{ij} + \frac{4}{\hat{\phi}} (\dot{\hat{\phi}} - \mathcal{L}_\xi \hat{\phi}) g_{ij} \right]. \quad (8)$$

The action (1) becomes

$$I_{\text{CS+V}} = \int d\lambda \int d^3x \sqrt{g} \hat{\phi}^4 \sqrt{R - 8 \frac{\nabla^2 \hat{\phi}}{\hat{\phi}}} \sqrt{\hat{T}}. \quad (9)$$

We now explain the rules by which we vary the action. A curve through the configuration space specifies a sequence $g_{ij}(\lambda)$ through Riem, combined with a λ -dependent conformal factor $\hat{\phi}(\lambda)$ and a λ -dependent coordinate transformation $\chi_i(\lambda)$ (for which $\xi_i = \partial \chi_i / \partial \lambda$). We identify ϕ and χ_i as gauge variables, so they need not reduce to the identity at the end points of the curve. Therefore, we extremize the action under fixed-end-point variations of g_{ij} and free-end-point variations of ϕ and χ_i .

We now illustrate free-end-point variation in a simple scale-invariant N -particle model [10]. The corrected coordinates are $\tilde{q}_{(i)} = a q_{(i)}$ and the corrected velocities are $\dot{\tilde{q}}_{(i)} = \dot{a} q_{(i)} + a \dot{q}_{(i)}$, $\mathcal{L} = \mathcal{L}(\tilde{q}_{(i)}, \dot{\tilde{q}}_{(i)})$. Here a is the scaling auxiliary and $q_{(i)}$ are the Cartesian coordinates of unit-mass point particles. The single auxiliary a matches the single scaling degeneracy of the $q_{(i)}$. It doubles the degeneracy. The habitual pairing of $q_{(i)}$ and a has several consequences, including full gauge invariance of all the theory’s equations, as we shall show elsewhere. More importantly, it imposes constraints. Indeed, the canonical

momenta, $p^{(i)} = \partial\mathcal{L}/\partial\dot{q}_i$ and $p^a = \partial\mathcal{L}/\partial\dot{a}$, are

$$\begin{aligned} p^{(i)} &= \frac{\partial\mathcal{L}}{\partial\dot{q}_{(i)}} \frac{\partial\dot{q}_{(i)}}{\partial\dot{q}_{(i)}} = \frac{\partial\mathcal{L}}{\partial\dot{q}_{(i)}} a, \\ p^a &= \frac{\partial\mathcal{L}}{\partial\dot{q}_{(i)}} \frac{\partial\dot{q}_{(i)}}{\partial\dot{a}} = \frac{\partial\mathcal{L}}{\partial\dot{q}_{(i)}} q_{(i)}. \end{aligned} \quad (10)$$

We therefore have the *primary constraint*

$$q_{(i)} p^{(i)} \equiv a p^a. \quad (11)$$

The variation $\int d\lambda (\delta\mathcal{L}/\delta a) \delta a$ is

$$\int d\lambda \left(\frac{dp^a}{d\lambda} - \frac{\partial\mathcal{L}}{\partial a} \right) \delta a + p^a (\delta a)_{\text{final}} - p^a (\delta a)_{\text{initial}}. \quad (12)$$

By the free-end-point rules, this must vanish for any δa . Starting with a general variation that vanishes at the end-points, we deduce the standard Euler-Lagrange equation for a : $dp^a/d\lambda = \partial\mathcal{L}/\partial a$. Then, choosing a variation that does not vanish at the initial end-point but does vanish at the final, and enforcing the Euler-Lagrange equations, we deduce that $p^a_{\text{initial}} = 0$. If we then choose a variation that does not vanish at the final end-point, we find that $p^a_{\text{final}} = 0$. As we can freely choose the final point along the curve, we deduce that $p^a = 0$. The Euler-Lagrange equation for a reduces to $\partial\mathcal{L}/\partial a = 0$. This is a key *consistency condition*. Hence, one can effectively minimize the action with respect to a and $\dot{a}$ independently, although the conditions really arise from a free-end-point variation of a . Finally, we get a *secondary constraint* $q_{(i)} p^{(i)} = 0$ from the primary constraint. These simple considerations capture all the novel features of our theory.

We may now proceed to apply this variational procedure to the conformalized BSW action (9). Variation with respect to $\dot{g}_{ij}$ defines the canonical momenta π^{ij} :

$$\pi^{ij} \equiv \frac{\delta I}{\delta \dot{g}_{ij}} = \Pi^{ij} + \frac{\langle \tilde{\Pi} \rangle}{3} \sqrt{g} g^{ij} (1 - \hat{\phi}^6), \quad (13)$$

where

$$\begin{aligned} \Pi^{ij} &= \frac{\sqrt{g}}{2N} G^{ijkl} \frac{d\hat{\phi}^4 g_{kl}}{d\lambda}, \\ N &\equiv \frac{1}{2} \sqrt{\frac{\hat{T}}{R - 8\frac{\nabla^2\phi}{\phi}}}, \end{aligned} \quad (14)$$

and $\Pi = \sqrt{g} \tilde{\Pi} = g_{ij} \Pi^{ij}$. Variation with respect to $\dot{\phi}$ gives

$$p_\phi \equiv \frac{\delta I}{\delta \dot{\phi}} = \frac{4}{\phi} (\Pi - \sqrt{g} \hat{\phi}^6 < \tilde{\Pi} >). \quad (15)$$

These definitions imply a primary constraint:

$$p_\phi = \frac{4}{\phi} (\pi - \sqrt{g} < p >), \quad (16)$$

where $\pi = \sqrt{g} p = g_{ij} \pi^{ij}$. Also, the rules of free-end-point variation imply that $p_\phi = 0$. Hence we find the secondary constraint

$$p = C, \quad (17)$$

where C is some spatial (ie λ -dependent) constant. This constraint is equivalent to the CMC condition imposed by York, obtained here as a genuine constraint and not a gauge-fixing condition (a similar result appeared earlier in [9]). As this condition designates a preferred time-function, it introduces a notion of global simultaneity.

The definitions of the momenta imply a quadratic primary identity, the analogue of the GR Hamiltonian constraint:

$$\begin{aligned} \pi^{ij} \pi_{ij} - \frac{\pi^2}{2} - \frac{C^2}{6} (1 - \hat{\phi}^6)^2 + \frac{C}{3} \sqrt{g} \pi (1 - \hat{\phi}^6) \\ = g \hat{\phi}^8 \left(R - 8 \frac{\nabla^2 \phi}{\phi} \right). \end{aligned} \quad (18)$$

Defining $\sigma^{ij} \equiv \pi^{ij} - \frac{1}{3} g^{ij} \pi$, and enforcing the CMC constraint (17), eqn. (18) takes the form

$$\sigma^{ij} \sigma_{ij} - \frac{\pi^2 \hat{\phi}^{12}}{6} - g \hat{\phi}^8 \left(R - 8 \frac{\nabla^2 \phi}{\phi} \right) = 0. \quad (19)$$

This identity holds true on any path in the configuration space. If we, however, switch to a Hamiltonian picture in which the canonical momenta become independent variables, it becomes, just as in standard GR, a real constraint. It is then clearly the Lichnerowicz-York equation [4].

The variation wrt ξ_i gives

$$\nabla_j \pi^{ij} = 0, \quad (20)$$

where we have imposed the CMC constraint (17); this relation is formally identical to the standard GR momentum constraint.

Variation with respect to ϕ and imposition of (17) leads to the consistency condition

$$\hat{\phi}^4 N \left(R - 7 \frac{\nabla^2 \phi}{\phi} \right) - \hat{\phi} \nabla^2 (\hat{\phi}^3 N) + \hat{\phi}^8 \frac{N p^2}{4} = \hat{\phi}^6 D, \quad (21)$$

where

$$D = \left\langle \hat{\phi}^4 N \left(R - 8 \frac{\nabla^2 \phi}{\phi} \right) + \frac{\hat{\phi}^8 N p^2}{4} \right\rangle. \quad (22)$$

One can show that D is the global average of the left-hand side of eqn. (21).

We now make a dimensional analysis. We take the metric, and by extension ϕ , to be dimensionless and give dimensions to the coordinates. We set the speed of light equal to one, so all coordinates have dimensions of length (l). Counting derivatives in their definitions, we find the

dimensions of other objects: $[R] = [\hat{R}] = [\hat{T}] = 1/l^2$. The lapse, N , is dimensionless. It follows that $[\pi^{ij}] = [\pi] = [p] = [C] = 1/l$. In the York scheme, one fixes an implicit length scale for the CWB manifold by specifying a numerical value for the dimensionful parameter C . That the scheme determines a physical 3-metric from an initially unphysical one reflects this fixing, as the manifold acquires a volume corresponding to the implicit length scale.

Our approach is complementary to this. We specify as initial data $g_{ij}, \dot{g}_{ij}$. Our procedure subjects them to considerable ‘gauge dressing’, but the initial volume V corresponding to the initial g_{ij} is the one thing that must not be changed. So we put the scale in explicitly through the canonical coordinate g_{ij} , not through a combination of the canonical momenta. In both cases, the specification of the numerical value of a single quantity with a length dimension fixes the numerical values of all other quantities with a length dimension. The time label is fixed only up to global reparametrization (as (21) is homogeneous in N), and, as always, the spatial coordinates are freely specifiable.

Our calculations have given us five equations—the CMC condition (17), the momentum constraint (20), and the consistency condition (21)—that we try to solve for the five gauge variables $(\hat{\phi}, \partial\hat{\phi}/\partial\lambda, \xi_i)$ given the initial $g_{ij}, \dot{g}_{ij}$. In accordance with our dimensional analysis the constant C , with its dimension l^{-1} , should emerge as an eigenvalue for this problem. There are many choices of $(g_{ij}, \dot{g}_{ij})$ for which we know solutions exist; e.g. take a CMC slice through a space-time that solves the Einstein equations, take the physical g_{ij} and $\dot{g}_{ij}$ given by the CMC foliation, and multiply by any VPCT. On the other hand, with a given arbitrary $(g_{ij}, \dot{g}_{ij})$, we have no existence theorem for solutions. This problem is very similar to the thin-sandwich problem in GR [7, 11].

If and when we can solve the equations, we can switch to the ‘corrected coordinates’ $\bar{g}_{ij}$. In this frame, $\hat{\phi} = 1$, and the consistency condition (21) reduces to

$$NR - \nabla^2 N + \frac{Np^2}{4} = D. \quad (23)$$

This relation is the CMC lapse-fixing equation of GR; satisfying it guarantees the propagation of the CMC constraint (17).

In contrast to the point of view of this article, York chose to retain the spacetime ontology and set up his powerful conformally-invariant method using invariance of the decoupling of the ADM Hamiltonian and momentum constraints as guiding principle. This led him to regard the Lichnerowicz–York equation as a gauge fixing

(see especially his comments near the end of the first paper in [4]). From our point of view, we rather choose to interpret the initial value problem as containing a genuine gauge invariance, which leads to gauge corrections both to the potential and to the metric velocities. From our point of view, we have derived GR, a prescription for solving its initial-value problem, and the condition for maintaining the CMC condition, all in a single package from our first principles. The transition to the Hamiltonian formulation and the implications of our results will be considered elsewhere.

We dedicate this paper to Jimmy York on the occasion of his 65th birthday. The authors thank him for discussions. EA also thanks Malcolm MacCallum for discussions and PPARC for financial support. EA, BF, BK, and NÓM thank the Barbour family for hospitality.

* Electronic address: eda@maths.qmul.ac.uk

† Electronic address: julian@platonica.com

‡ Electronic address: bzf@physics.umd.edu

§ Electronic address: bk@physics.ucc.ie

¶ Electronic address: niall@ucc.ie

- [1] R. Arnowitt, S. Deser, and C. Misner in *Gravitation: an Introduction to Current Research*, ed. L. Witten, (Wiley, New York, 1962).
- [2] P. A. M. Dirac, *Lectures on Quantum Mechanics* (Yeshiva University, New York).
- [3] J. W. York, Ann. Inst. Henri Poincaré **21**, 319 (1974); A. E. Fischer and V. Moncrief, Gen. Rel. Grav. **28**, 221 (1996).
- [4] J. W. York, Phys. Rev. Lett. **26**, 1656 (1971); J. W. York, Phys. Rev. Lett. **28**, 1082 (1972); J. W. York, J. Math. Phys. **14**, 456 (1973).
- [5] C. W. Misner, K. Thorne and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
- [6] A. Lichnerowicz, J. Math. Pures Appl. **23**, 37 (1944).
- [7] R. F. Baierlein, D. Sharp and J. A. Wheeler, Phys. Rev. Lett. **126**, 1864 (1962).
- [8] J. Barbour, B. Z. Foster and N. O’Murchadha, Class. Quant. Grav. **19**, 3217 (2002) E. Anderson, Phys. Rev. D **68**, 104001 (2003) E. Anderson, PhD Thesis: *Geometrodynamics: Spacetime or Space?*, University of London (2004).
- [9] J. Barbour and N. O. Murchadha, arXiv:gr-qc/9911071; E. Anderson, J. Barbour, B. Foster and N. O’Murchadha, Class. Quant. Grav. **20**, 1571 (2003) B. Kelleher, Class. Quant. Grav. **21**, 2623 (2004).
- [10] J. Barbour, Class. Quant. Grav. **20**, 1543 (2003)
- [11] E. P. Belasco and H. C. Ohanian, J. Math. Phys. **10**, 1053 (1969); R. Bartnik and G. Fodor, Phys. Rev. D **48**, 3596 (1993) D. Giulini, J. Math. Phys. **40**, 2470 (1999)